

The 5 March 1979 soft gamma-ray repeater: a fresh outlook

G.V. Vlasov *

February 1, 2008

Abstract

An attempt is made to explain the 8-second periodicity of the 5th March 1979 soft gamma-ray repeater as a consequence of the elastic wave precession in the rigid crust of the rotating neutron star.

The soft gamma-repeaters (SGR) as well as the ordinary gamma-ray bursts (GRB) are observed for almost thirty years. While the origin of GRB still remains mysterious [1], the SGR are associated with the neutron stars [2, 1, 3, 4, 5], particularly, the most incomparable event enrolled on the 5th of March 1979 [6] from SGR 0526-66, which arouse most inquiry and discussion. Its source is believed to be a neutron star, the remainder of supernova N 49 [7, 8, 9, 1]. The radiation of that SGR is modulated at definite frequencies [10, 11], whereas the most intense 8-second peak is usually associated with period P of star rotation [12, 13], the origin of the 4-second [10, 11] and 6-second peaks [10] remains unknown. The rigid crust radial vibration is suggested to take place in such events [14, 12], yet, those periodicities cannot be provoked by vibration itself, since the relevant characteristic frequencies ($\nu \sim 1ms^{-1}$) are too high (other examples of SGR with periods $P \sim 1s$ are also known [15]). We guess the nature of those "attendant" frequencies, as well as the main period, can be explained as soon as we appeal to an interesting phenomenon neglected so far. It is the inertia of elastic waves in a rotating

*E-mail: vs@itp.ac.ru

rigid body, first discovered a century ago [16]. The investigation of a new gyroscope [17, 18] was nourished from this effect.

Suppose that the elastic vibrations are excited in the solid crust of a neutron star. The spectrum of vibrations is a discrete set of frequencies ν_k denoted by integer k . If the body is put into rotation at frequency $\omega = 2\pi/P$, then, for each k -mode, there exists a reference frame within which the relevant standing wave is possible. But this reference frame rotates itself at definite angular velocity [18]

$$\Omega = \omega(1 - \chi) \quad \chi = \frac{kA_n^2}{1 + n(n+1)A_n^2} \quad 2 \leq k \leq n \quad 0 < \chi < 1 \quad (1)$$

where parameter A_n is a function of the number of waves n along the meridian and the Poisson coefficient μ [19]:

$$A_n = \frac{-2(1 + \mu) + \frac{1}{2}(1 - \mu)\lambda}{(1 + \mu)n(n+1)} \quad (2)$$

where λ is determined by equation

$$(1 - \mu)\lambda^2 - 2\nu[1 + 3\mu + n(n+1)] + 4(1 + \mu)(n^2 + n - 2) = 0 \quad (3)$$

By the way, the auxiliary modes (marked by tilde) are also possible here. For the body at rest all Ω are the same and equal to ω . In other words, one finds the vibration modes traveling, each at definite period $T = 2\pi/\Omega$ and not equal to P . In addition, the larger k , the lesser T , so that $\Omega \rightarrow \omega$ when $k \rightarrow \infty$. This phenomenon is known as inertia of the elastic waves, for the elastic vibrations behave as if they were a mass involved into precession.

For the matter of the neutron star crust [20] with $\mu \simeq 1/2$ the main and auxiliary period of the simplest dipole mode ($n = k = 2$), obtained in the frames of theory [18] by Eqs. (1)-(3), is estimated as $1.28P$ and $1.13P$ respectively.

However, for $n = 2$ and $k = 4$ we come to a system of fractional connectedness. In the light of crustal plate tectonics during gamma-bursts [20, 21, 22], the crust cracking and splitting in two halves is the simplest event of conceivable situations. Indeed, the case $k = 2n$, corresponds to the two vibrating semi-spheres. The formula (1) then is reduced to [23]

$$\chi = \frac{-2 - \mu \pm \sqrt{(2 + \mu)^2 + 4(1 - \mu)k^2}}{2(1 - \mu)k^2} \simeq \frac{-5 \pm \sqrt{25 + 8k^2}}{2k^2} \quad (4)$$

and it gives $T = 1.30P$ and $\tilde{T} = 0.65P$ at $k = 4$; if $P = 6s$, then $T = 7.8s$ and $\tilde{T} = 3.9s$. Of course, we may vary k to obtain various ratio $T/\tilde{T}$.

If we rely on the theory of relationship between gamma-radiation and star crust vibration, as it has been claimed once [14], the observer will reveal radiation modulated at frequencies Ω determined by the formulae (1), (4) and in agreement with the observations [10]. Note that the frequency ω itself may not be discovered at all. Were there no inertia of elastic waves in a rigid crust, the observer would detect only the single frequency ω instead of a set $\{\Omega\}$.

Whereas, the real process includes other factors not discussed in this simple consideration, it is clear that the most intensive 8-second peak scarcely coincides with the period of star rotation. Since the lowest modes are the most powerful, the 8-second period probably corresponds to the main dipole mode of a vibrating sphere or two semi-spheres. Although we have not taken into account all the details, for instance the general relativistic effects, the role of finite ratio Ω/ν_k and thickness of the crust, the qualitative result remains the same anyhow. The precession of elastic vibrations in the solid crust of a rotating neutron star leads to a qualitatively new effect and the radiation of SGR shall be modulated at several frequencies instead of a single one.

References

- [1] Ch. Kouveliotou *et al*, Nature **368**, 125 (1994).
- [2] S.R. Kulkarni and D.A. Frail, Nature **365**, 33 (1993).
- [3] T. Murakami *et al*, Nature **368**, 127 (1994).
- [4] S.R. Kulkarni *et al*, Nature **368**, 129 (1994).
- [5] K.S. Cheng and Z.G.Dai, Phys. Rev. Lett. **80**, 18 (1998).
- [6] E.P. Mazets *et al*, Nature **282**, 587 (1979).
- [7] D.J. Helfand and K.S. Long, Nature **282**, 589 (1979).
- [8] W.D. Evans *et al*, Astrophys. J. **237**, L7 (1980).

- [9] T.L. Cline *et al*, *Astrophys. J.* **255**, L45 (1982).
- [10] C. Barat *et al*, *Astron. Astrophys.* **79**, L24 (1979).
- [11] J. Terrell *et al*, *Nature* **285**, 383 (1980).
- [12] D. Kazanas, *Nature* **331**, 320 (1988).
- [13] Ch. Alcock, E. Farthi, and A. Olinto, *Phys. Rev. Lett.* **57**, 2088 (1986).
- [14] R. Ramaty *et al*, *Nature* **287**, 122 (1980).
- [15] Ch. Kouveliotou *et al*, *astro-ph/9809140*.
- [16] G.H. Bryan, *Proc. Cam. Philos. Soc. Math. Phys. Sci.* **7**, 101 (1890).
- [17] W.B. Scott, *Aviat. Week & Space Technol.* **117**, n.17, 64 (1982).
- [18] V.F. Zhuravlev, *Mech. Solids* **28**, n.3, 3 (1993).
- [19] V.F. Zhuravlev, *Mech. Solids* **21**, n.6, 87 (1986).
- [20] M. Ruderman, *Astrophys. J.* **366**, 261 (1991).
- [21] M. Ruderman, *Astrophys. J.* **382**, 587 (1991).
- [22] B. Link, L.M. Franco and R.I. Epstein, *astro-ph/9805115*.
- [23] V.F. Zhuravlev and A.L. Popov, *Mech. Solids* **20** n.1, 138 (1985).